arXiv:math/0312150v1 [math.CO] 8 Dec 2003

The Pseudo Cosine Sequences of a Distance-Regular Graph

Arlene A. Pascasio Paul M. Terwilliger

July 11,2003

Abstract

Let Γ denote a distance-regular graph with diameter $D \geq 3$, valency k, and intersection numbers a_i, b_i, c_i. By a *pseudo cosine sequence* of Γ we mean a sequence of real numbers $\sigma_0, \sigma_1, \ldots, \sigma_D$ such that $\sigma_0 = 1$ and $c_i\sigma_{i-1} + a_i\sigma_i + b_i\sigma_{i+1} = k\sigma_1\sigma_i$ for $0 \leq i \leq D-1$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote pseudo cosine sequences of Γ. We say this pair of sequences is *tight* whenever $\sigma_0\rho_0, \sigma_1\rho_1, \ldots, \sigma_D\rho_D$ is a pseudo cosine sequence of Γ. In this paper, we determine all the tight pairs of pseudo cosine sequences of Γ.

AMS Classification: 05E50
Keywords: distance-regular graph, association scheme, cosine sequence, pseudo cosine sequence.

1 Introduction

Let $\Gamma = (X, R)$ denote a distance-regular graph with diameter $D \geq 3$, valency k, and intersection numbers a_i, b_i, c_i (see Section 2 for formal definitions). Let θ and θ' denote eigenvalues of Γ other than k. In [9], Jurišić, Koolen and Terwilliger showed

$$\left(\theta + \frac{k}{a_1+1}\right)\left(\theta' + \frac{k}{a_1+1}\right) \geq -\frac{ka_1b_1}{(a_1+1)^2}. \tag{1}$$

The case of equality in (1) has received a lot of attention. This case has been characterized using the intersection numbers [9, Theorem 10.2], the 1-homogeneous property [9, Theorem 11.7(i)], and the local structure [9, Theorem 12.6]. See also [3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. In [14], Pascasio characterized equality in (1) using cosine sequences. The idea is as follows. Let θ denote an eigenvalue of Γ. Recall the *cosine sequence* for θ is the sequence of real numbers $\sigma_0, \sigma_1, \ldots, \sigma_D$ such that $\sigma_0 = 1$ and $c_i\sigma_{i-1} + a_i\sigma_i + b_i\sigma_{i+1} = \theta\sigma_i$ for $0 \leq i \leq D$, where σ_{-1} and σ_{D+1} are indeterminate. Let θ and θ' denote eigenvalues of Γ other than k. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ (resp. $\rho_0, \rho_1, \ldots, \rho_D$) denote the cosine sequence for θ (resp. θ'). Then by [14, Theorem 3.4], θ, θ' satisfy (1) with equality if and only if $\sigma_0\rho_0, \sigma_1\rho_1, \ldots, \sigma_D\rho_D$ is a cosine sequence.

The concept of a *pseudo cosine sequence* was recently introduced [10, 19]. This concept is defined as follows. Let θ denote a real number. By the *pseudo cosine sequence* for θ

we mean the sequence of real numbers $\sigma_0, \sigma_1, \ldots, \sigma_D$ such that $\sigma_0 = 1$ and $c_i\sigma_{i-1} + a_i\sigma_i + b_i\sigma_{i+1} = \theta\sigma_i$ for $0 \leq i \leq D-1$, where σ_{-1} is indeterminate. Pseudo cosine sequences have been used to describe certain modules for the subconstituent algebra [19]. They also arise in connection with the pseudo 1-homogeneous property [10]. Given our comments in the previous paragraph it is natural to consider the following situation. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote pseudo cosine sequences of Γ. We say this pair of sequences is *tight* whenever $\sigma_0\rho_0, \sigma_1\rho_1, \ldots, \sigma_D\rho_D$ is a pseudo cosine sequence of Γ.

In this paper we find all the tight pairs of pseudo cosine sequences of Γ. We break the argument into the following cases: (i) $a_i = 0$ for $0 \leq i \leq D-1$; (ii) $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$; (iii) $a_1 = 0$ and there exists i $(2 \leq i \leq D-2)$ such that $a_i \neq 0$; (iv) $a_1 \neq 0$. Our results for these cases are given in Theorem 5.3, Theorem 9.4, Theorem 10.1, and Theorem 14.2 respectively.

Our methods in this paper are purely algebraic. Consequently our main results probably hold for P-polynomial character algebras. We will pursue this in a future paper.

2 Preliminaries

In this section, we review some definitions and basic concepts. For more background information, the reader may refer to the books of Bannai and Ito [1], or Brouwer, Cohen and Neumaier [2].

Let $\Gamma = (X, R)$ denote a finite, undirected, connected graph without loops or multiple edges, with vertex set X, edge set R, path-length distance function ∂ and diameter $D := \max\ \{\partial(x,y) | x, y \in X\}$. Let k denote a nonnegative integer. We say Γ is *regular with valency* k whenever each vertex of Γ is adjacent to exactly k distinct vertices of Γ. We say Γ is *distance-regular* whenever for all integers h, i, j $(0 \leq h, i, j \leq D)$ and for all $x, y \in X$ with $\partial(x,y) = h$, the number

$$p^h_{ij} := |\{z \in X |\ \partial(x,z) = i,\ \partial(y,z) = j\}|$$

is independent of x and y. The integers p^h_{ij} are called the *intersection numbers* for Γ. We abbreviate $a_i := p^i_{1i}$ $(0 \leq i \leq D)$, $b_i := p^i_{1i+1}$ $(0 \leq i \leq D-1)$, $c_i := p^i_{1i-1}$ $(1 \leq i \leq D)$, and $k_i := p^0_{ii}$ $(0 \leq i \leq D)$. We observe $a_0 = 0$, $c_1 = 1$, and $k_0 = 1$. For notational convenience we define $c_0 := 0$ and $b_D := 0$. For the rest of this paper we assume Γ is distance-regular with diameter $D \geq 3$.

We have a few comments. The graph Γ is regular with valency $k = k_1 = b_0$. Moreover

$$c_i + a_i + b_i = k \qquad (0 \leq i \leq D). \tag{2}$$

It is known, by [1, Chapter 3, Proposition 1.2]

$$k_i = \frac{b_0 b_1 \cdots b_{i-1}}{c_1 c_2 \cdots c_i} \qquad (0 \leq i \leq D). \tag{3}$$

We remark $c_{i-1} \leq c_i$ and $b_{i-1} \geq b_i$ for $1 \leq i \leq D$ [2, Proposition 4.1.6].

We recall the Bose-Mesner algebra of Γ. Let $\mathrm{Mat}_{\mathrm{X}}(\mathbb{C})$ denote the $\mathbb{C}$-algebra consisting of the matrices which have rows and columns indexed by X and entries in $\mathbb{C}$. For $0 \leq i \leq D$ let A_i denote the matrix in $\mathrm{Mat}_{\mathrm{X}}(\mathbb{C})$ with entries

$$(A_i)_{xy} = \begin{cases} 1, & \text{if } \partial(x,y) = i \\ 0, & \text{if } \partial(x,y) \neq i \end{cases} \qquad (x, y \in X).$$

We call A_i the *ith distance matrix* of Γ. We have $A_0 = I$, $\sum_{i=0}^{D} A_i = J$, $A_i^t = A_i$ $(0 \leq i \leq D)$, $A_i A_j = \sum_{h=0}^{D} p_{ij}^h A_h$ $(0 \leq i, j \leq D)$ where J denotes the all 1's matrix. The matrices $A_0, A_1, \ldots, A_D$ form a basis for a commutative semi-simple $\mathbb{C}$-algebra M, called the *Bose-Mesner algebra of* Γ. By [1, Section 2.3] M has second basis $E_0, E_1, \ldots, E_D$ such that $E_0 = |X|^{-1}J$, $\sum_{i=0}^{D} E_i = I$, $E_i^t = E_i$ $(0 \leq i \leq D)$, $E_i E_j = \delta_{ij} E_i$ $(0 \leq i, j \leq D)$. The $E_0, E_1, \ldots, E_D$ are called the *primitive idempotents* of Γ. We call E_0 the *trivial* idempotent of Γ.

We set $A := A_1$ and call this the *adjacency matrix* of Γ. Let $\theta_0, \theta_1, \ldots, \theta_D$ denote the complex scalars such that $A = \sum_{i=0}^{D} \theta_i E_i$. It is known $\theta_0 = k$, and that $\theta_0, \theta_1, \ldots, \theta_D$ are distinct real numbers [1, Chapter 3, Theorem 1.3]. We refer to θ_i as the *eigenvalue of* Γ *associated with* E_i $(0 \leq i \leq D)$. We call θ_0 the *trivial eigenvalue of* Γ. For each integer i $(0 \leq i \leq D)$, let m_i denote the rank of E_i. We refer to m_i as the *multiplicity* of E_i (or θ_i).

Let θ denote an eigenvalue of Γ, let E denote the associated primitive idempotent, and let m denote the multiplicity of E. By [1, Section 2.3] there exists a sequence of real numbers $\sigma_0, \sigma_1, \ldots, \sigma_D$ such that

$$E = |X|^{-1} m \sum_{i=0}^{D} \sigma_i A_i.$$

It follows from [1, Chapter 2, Proposition 3.3 (iii)] that $\sigma_0 = 1$. We call $\sigma_0, \sigma_1, \ldots, \sigma_D$ the *cosine sequence of* Γ *associated with* θ. We shall often abbreviate σ_1 by σ. Let $\sigma_0, \sigma_1, \ldots, \sigma_D, \theta$ denote real numbers. By [2, Proposition 4.1.1] the following are equivalent: (i) θ is an eigenvalue of Γ and $\sigma_0, \sigma_1, \ldots, \sigma_D$ is the corresponding cosine sequence; (ii) $\sigma_0 = 1$ and $c_i \sigma_{i-1} + a_i \sigma_i + b_i \sigma_{i+1} = \theta \sigma_i$ $(0 \leq i \leq D)$, where σ_{-1} and σ_{D+1} are indeterminates.

We end this section with a comment.

Lemma 2.1. [2, Proposition 5.5.1] *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$. *Suppose* $a_1 \neq 0$. *Then* $a_i \neq 0$ *for* $1 \leq i \leq D-1$.

3 Pseudo cosine sequences

We now recall a sequence of real numbers which generalizes the cosine sequence.

Definition 3.1. *[10, 19]* Let Γ denote a distance-regular graph with diameter $D \geq 3$. For $\theta \in \mathbb{R}$, by the *pseudo cosine sequence for θ* we mean the sequence of real numbers $\sigma_0, \sigma_1, \ldots, \sigma_D$ such that $\sigma_0 = 1$ and

$$c_i \sigma_{i-1} + a_i \sigma_i + b_i \sigma_{i+1} = \theta \sigma_i \quad (0 \leq i \leq D-1), \tag{4}$$

where σ_{-1} is an indeterminate. For notational convenience we abbreviate $\sigma := \sigma_1$. We remark θ is determined by its pseudo cosine sequence; indeed $\theta = k\sigma$, where k denotes the valency of Γ.

We have some remarks on Definition 3.1. Let Γ denote a distance-regular graph with diameter $D \geq 3$. Pick $\theta \in \mathbb{R}$ and let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the corresponding pseudo cosine sequence. Then θ is an eigenvalue of Γ if and only if $c_D \sigma_{D-1} + a_D \sigma_D = \theta \sigma_D$. In this case $\sigma_0, \sigma_1, \ldots, \sigma_D$ is the cosine sequence for θ. Now let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence for $\theta = k$, where k is the valency of Γ. Comparing (2), (4) we find $\sigma_i = 1$ for $0 \leq i \leq D$. By the *trivial* pseudo cosine sequence we mean the pseudo cosine sequence for k. This sequence is a cosine sequence since k is an eigenvalue of Γ.

Lemma 3.2. (Christoffel-Darboux) *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote pseudo cosine sequences of Γ. Then*

$$(\sigma - \rho) \sum_{h=0}^{i} k_h \sigma_h \rho_h = \frac{b_1 b_2 \cdots b_i}{c_1 c_2 \cdots c_i} (\sigma_{i+1} \rho_i - \sigma_i \rho_{i+1}) \qquad (0 \leq i \leq D-1). \tag{5}$$

Proof. For $0 \leq h \leq i$ we have

$$\begin{aligned} \sigma \sigma_h k &= c_h \sigma_{h-1} + a_h \sigma_h + b_h \sigma_{h+1}, & (6) \\ \rho \rho_h k &= c_h \rho_{h-1} + a_h \rho_h + b_h \rho_{h+1}. & (7) \end{aligned}$$

Evaluating the left-hand side of (5) using (6), (7), and simplifying the result using (3) we routinely obtain the right-hand side of (5). $\square$

Lemma 3.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote real numbers. Then the following are equivalent.*

(i) $\sigma_0, \sigma_1, \ldots, \sigma_D$ *is a pseudo cosine sequence.*

(ii) $\sigma_0 = 1$ *and*

$$c_i(\sigma_{i-1} - \sigma_i) - b_i(\sigma_i - \sigma_{i+1}) = k(\sigma - 1)\sigma_i \qquad (1 \leq i \leq D-1). \tag{8}$$

(iii) $\sigma_0 = 1$ *and*

$$(\sigma - 1) \sum_{h=0}^{i} k_h \sigma_h = \frac{b_1 b_2 \cdots b_i}{c_1 c_2 \cdots c_i} (\sigma_{i+1} - \sigma_i) \qquad (0 \leq i \leq D-1). \tag{9}$$

Proof. (i)$\Longrightarrow$(iii) To show (9), in Lemma 3.2 let $\rho_0, \rho_1, \ldots, \rho_D$ denote the pseudo cosine sequence for k.
(iii)$\Longrightarrow$(ii) We show (8) holds. Pick an integer i $(1 \leq i \leq D-1)$. By (9) (with i replaced by $i-1$) we have

$$(\sigma - 1)\sum_{h=0}^{i-1} k_h \sigma_h = \frac{b_1 b_2 \cdots b_{i-1}}{c_1 c_2 \cdots c_{i-1}}(\sigma_i - \sigma_{i-1}). \tag{10}$$

Subtracting equation (10) from equation (9) and eliminating k_i from the result using (3) we get (8) as desired.
(ii)$\Longrightarrow$(i) We show

$$c_i \sigma_{i-1} + a_i \sigma_i + b_i \sigma_{i+1} = k\sigma\sigma_i \tag{11}$$

for $0 \leq i \leq D-1$. Clearly (11) holds for $i = 0$. To show (11) holds for $1 \leq i \leq D-1$ simplify (8) using (2). We now have (11) for $0 \leq i \leq D-1$. Applying Definition 3.1 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ is the pseudo cosine sequence for $k\sigma$. □

We mention another characterization of the pseudo cosine sequences.

Lemma 3.4. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote real numbers. Then the following are equivalent.*

(i) $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a pseudo cosine sequence.

(ii) $\sigma_0 = 1$ and

$$b_i(\sigma_{i-1} - \sigma_{i+1}) = k(\sigma_{i-1} - \sigma\sigma_i) - a_i(\sigma_{i-1} - \sigma_i) \qquad (1 \leq i \leq D-1). \tag{12}$$

(iii) $\sigma_0 = 1$ and

$$c_i(\sigma_{i+1} - \sigma_{i-1}) = k(\sigma_{i+1} - \sigma\sigma_i) - a_i(\sigma_{i+1} - \sigma_i) \qquad (1 \leq i \leq D-1). \tag{13}$$

Proof. (i)$\Longrightarrow$(ii) Lemma 3.3(i) holds. Applying that lemma and eliminating c_i in (8) using (2) we get (12).
(ii)$\Longrightarrow$(iii) Eliminating b_i in (12) using (2) we get (13).
(iii)$\Longrightarrow$(i) We show (8) holds. To do this we eliminate a_i in (13) using (2). It follows $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a pseudo cosine sequence in view of Lemma 3.3(i),(ii). □

We finish this section with some results which we will find useful later in the paper.

Lemma 3.5. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a pseudo cosine sequence of Γ. Then at least one of σ_i, σ_{i+1} is nonzero for $0 \leq i \leq D-1$.*

Proof. Assume there exists an integer i $(0 \leq i \leq D-1)$ such that $\sigma_i = 0$ and $\sigma_{i+1} = 0$. Recall $\sigma_0 = 1$ so $i \geq 1$. Without loss of generality we may assume $\sigma_{i-1} \neq 0$. This is inconsistent with (4) and the fact that $c_i \neq 0$. The result follows. □

Lemma 3.6. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a nontrivial pseudo cosine sequence of Γ. Then for $1 \leq i \leq D-1$, at least one of $\sigma_{i-1} - \sigma_i$, $\sigma_i - \sigma_{i+1}$ is nonzero.*

Proof. Suppose there exists an integer i $(1 \leq i \leq D-1)$ such that each of $\sigma_{i-1} - \sigma_i$, $\sigma_i - \sigma_{i+1}$ is zero. Then $\sigma_{i-1}, \sigma_i, \sigma_{i+1}$ coincide and this common value is nonzero by Lemma 3.5. Evaluating (8) using this we find $\sigma = 1$. Now $\sigma_0, \sigma_1, \ldots, \sigma_D$ is the trivial pseudo cosine sequence, a contradiction. ☐

Lemma 3.7. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence of Γ associated with $-k$. Suppose there exists an integer i $(1 \leq i \leq D-2)$ such that $\sigma_{i-1} + \sigma_i = 0$ and $\sigma_{i+1} + \sigma_{i+2} = 0$. Then $a_i = 0$, $a_{i+1} = 0$, and $\sigma_i + \sigma_{i+1} = 0$.*

Proof. Observe $\sigma_i \neq 0$; otherwise each of σ_{i-1}, σ_i is zero, contradicting Lemma 3.5. Also $\sigma_{i+1} \neq 0$; otherwise each of $\sigma_{i+1}, \sigma_{i+2}$ is zero, contradicting Lemma 3.5. Applying (4) at i and $i+1$ with $\theta = -k$ we find both $b_i \sigma_{i+1} = \sigma_i(c_i - a_i - k)$ and $c_{i+1}\sigma_i = \sigma_{i+1}(b_{i+1} - a_{i+1} - k)$. Combining these two equations and using (2) we routinely obtain

$$(k + a_i - c_i)a_{i+1} = -a_i c_{i+1}. \tag{14}$$

In (14) the left-hand side is nonnegative and the right-hand side is nonpositive. Therefore both sides are zero. It follows each of a_i, a_{i+1} is zero. Setting $a_i = 0$ in the equation $b_i \sigma_{i+1} = \sigma_i(c_i - a_i - k)$ we routinely obtain $\sigma_i + \sigma_{i+1} = 0$. ☐

4 The tight relation

Definition 4.1. Let Γ denote a distance-regular graph with diameter $D \geq 3$. We introduce a binary symmetric relation on the set of all pseudo cosine sequences of Γ. We call this the *tight* relation. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote pseudo cosine sequences of Γ. We say this pair of sequences is *tight* whenever $\sigma_0\rho_0, \sigma_1\rho_1, \ldots, \sigma_D\rho_D$ is a pseudo cosine sequence of Γ.

Definition 4.2. Let Γ denote a distance-regular graph with diameter $D \geq 3$. We define a binary symmetric relation on $\mathbb{R}$ which we call the *tight* relation. For $\theta, \theta' \in \mathbb{R}$, we say the pair θ, θ' is *tight* whenever the corresponding pseudo cosine sequences form a tight pair.

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. We give an example of a tight pair of pseudo cosine sequences. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence for k and recall $\sigma_i = 1$ for $0 \leq i \leq D$. Let $\rho_0, \rho_1, \ldots, \rho_D$ denote any pseudo cosine sequence. Of course $\sigma_i\rho_i = \rho_i$ for $0 \leq i \leq D$. Applying Definition 4.1 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Consequently we have the following.

Example 4.3. Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Then for all $\theta \in \mathbb{R}$ the pair θ, k is tight.

Lemma 4.4. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Let θ, θ' denote real numbers other than k, and assume θ, θ' form a tight pair. Then*

$$\left(\theta + \frac{k}{a_1 + 1}\right)\left(\theta' + \frac{k}{a_1 + 1}\right) = -\frac{k a_1 b_1}{(a_1 + 1)^2}. \tag{15}$$

Proof. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ (resp. $\rho_0, \rho_1, \ldots, \rho_D$) denote the pseudo cosine sequence for θ (resp. θ'). Define $\gamma_i := \sigma_i \rho_i$ for $0 \leq i \leq D$. Observe $\gamma_0, \gamma_1, \ldots, \gamma_D$ is the pseudo cosine sequence for ψ, where $\psi := k\sigma\rho$. Setting $i = 1$ in (4) and using $\theta = k\sigma$ we find

$$1 + a_1\sigma + b_1\sigma_2 = k\sigma^2. \tag{16}$$

Similarly

$$\begin{aligned} 1 + a_1\rho + b_1\rho_2 &= k\rho^2, & (17) \\ 1 + a_1\sigma\rho + b_1\sigma_2\rho_2 &= k\sigma^2\rho^2. & (18) \end{aligned}$$

To obtain (15), first solve (16) and (17) for σ_2 and ρ_2 respectively. Then evaluate (18) using this and simplify the result using $\theta = k\sigma$, $\theta' = k\rho$ and (2). $\square$

In this paper, we will find all the tight pairs. We will break down the argument into the following cases (i) $a_i = 0$ for $0 \leq i \leq D-1$; (ii) $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$; (iii) $a_1 = 0$ and there exists i $(2 \leq i \leq D-2)$ such that $a_i \neq 0$; (iv) $a_1 \neq 0$.

5 The case $a_i = 0$ $(0 \leq i \leq D-1)$

Let Γ denote a distance-regular graph with diameter $D \geq 3$. Recall Γ is *bipartite* whenever $a_i = 0$ for $0 \leq i \leq D$. We say Γ is *almost bipartite* whenever $a_D \neq 0$ and $a_i = 0$ for $0 \leq i \leq D-1$.

Lemma 5.1. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Assume Γ is bipartite or almost bipartite. Given $\theta \in \mathbb{R}$, let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence for θ and let $\rho_0, \rho_1, \ldots, \rho_D$ denote the pseudo cosine sequence for $-\theta$. Then $\rho_i = (-1)^i \sigma_i$ for $0 \leq i \leq D$.*

Proof. Observe the desired result holds for $i = 0$ and $i = 1$ since $\sigma_0 = 1 = \rho_0$ and $\sigma = \theta k^{-1} = -\rho$. To show the result holds for $2 \leq i \leq D$ we use induction and the fact that $c_{i-1}\sigma_{i-2} + b_{i-1}\sigma_i = \theta\sigma_{i-1}$ and $c_{i-1}\rho_{i-2} + b_{i-1}\rho_i = -\theta\rho_{i-1}$. $\square$

Corollary 5.2. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Assume Γ is bipartite or almost bipartite. Let $\rho_0, \rho_1, \ldots, \rho_D$ denote the pseudo cosine sequence for $-k$. Then $\rho_i = (-1)^i$ for $0 \leq i \leq D$.*

Proof. Referring to Lemma 5.1, set $\theta = k$ and recall $\sigma_i = 1$ for $0 \leq i \leq D$. $\square$

Theorem 5.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Assume Γ is bipartite or almost bipartite.*

(i) The pair θ, k is tight for all $\theta \in \mathbb{R}$.

(ii) The pair $\theta, -k$ is tight for all $\theta \in \mathbb{R}$.

(iii) Γ has no further tight pairs.

Proof. (i) This is from Example 4.3.
(ii) Let $\sigma_0, \sigma_1, \dots, \sigma_D$ denote the pseudo cosine sequence for θ, and let $\rho_0, \rho_1, \dots, \rho_D$ denote the pseudo cosine sequence for $-k$. We show the sequences $\sigma_0, \sigma_1, \dots, \sigma_D$ and $\rho_0, \rho_1, \dots, \rho_D$ form a tight pair. For $0 \leq i \leq D$ define $\gamma_i := \sigma_i \rho_i$. Recall $\rho_i = (-1)^i$ for $0 \leq i \leq D$ by Corollary 5.2 so $\gamma_i = (-1)^i \sigma_i$. Observe $\gamma_0, \gamma_1, \dots, \gamma_D$ is the pseudo cosine sequence for $-\theta$ by Lemma 5.1. We have now shown the sequences $\sigma_0, \sigma_1, \dots, \sigma_D$ and $\rho_0, \rho_1, \dots, \rho_D$ form a tight pair. It follows the pair $\theta, -k$ is tight.
(iii) For $\theta \in \mathbb{R}$, $\theta' \in \mathbb{R}$, suppose θ, θ' is a tight pair but $\theta \neq k$, $\theta' \neq k$. We show $\theta = -k$ or $\theta' = -k$. Applying Lemma 4.4 we find θ, θ' satisfy (15). By this and since $a_1 = 0$ we find $(\theta + k)(\theta' + k) = 0$. Now $\theta = -k$ or $\theta' = -k$. The result follows. □

6 Tight pairs θ, θ' with $\theta = \theta'$.

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Let θ, θ' denote real numbers other than k, and assume the pair θ, θ' is tight. Conceivably $\theta = \theta'$; we consider when this occurs. First suppose Γ is bipartite or almost bipartite. ¿From Lemma 5.3 we find $\theta = \theta'$ if and only if each of θ, θ' is equal to $-k$. We now treat the case in which Γ is neither bipartite nor almost bipartite.

Lemma 6.1. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Assume Γ is neither bipartite nor almost bipartite. Let θ, θ' denote real numbers other than k, and assume the pair θ, θ' is tight. Then $\theta' \neq \theta$.*

Proof. Applying Lemma 4.4 we find θ, θ' satisfies (15). Suppose $\theta = \theta'$. Then the left-hand side of (15) is a square so it is nonnegative. The right-hand side of (15) is nonpositive so both sides of (15) are equal to 0. Now $a_1 = 0$. Moreover θ and θ' are equal to $-k$. Let $\sigma_0, \sigma_1, \dots, \sigma_D$ denote the pseudo cosine sequence for θ. Since the pair θ, θ is tight we find $\sigma_0^2, \sigma_1^2, \dots, \sigma_D^2$ is a pseudo cosine sequence. This is a pseudo cosine sequence for ψ where $\psi := k\sigma^2$. Recall $\sigma = \theta/k$ and we mentioned $\theta = -k$ so $\sigma = -1$. Apparently $\psi = k$ so $\sigma_i^2 = 1$ for $0 \leq i \leq D$. We assume Γ is neither bipartite nor almost bipartite. Therefore there exists an integer r $(0 \leq r \leq D - 1)$ such that $a_r \neq 0$. Assume we have picked the minimal such r. Observe $r \geq 2$ since $a_0 = 0$ and $a_1 = 0$. For $0 \leq i < r$ we apply (4) with $\theta = -k$ and $a_i = 0$ to obtain $\sigma_j = (-1)^j$ for $0 \leq j \leq r$. We now apply (4) with $i = r$ and $\theta = -k$. Simplifying the result using $\sigma_{r-1} = (-1)^{r-1}$, $\sigma_r = (-1)^r$ and (12) we obtain

$$1 - (-1)^{r+1}\sigma_{r+1} = -2a_r/b_r. \tag{19}$$

¿From our above comments $\sigma_{r+1}^2 = 1$ so $\sigma_{r+1} = 1$ or $\sigma_{r+1} = -1$. It follows the left-hand side of (19) is nonnegative. By construction $a_r > 0$ so the right-hand side of (19) is negative. We now have a contradiction. We conclude $\theta \neq \theta'$. □

7 The auxiliary parameter

We now introduce a parameter which will help us describe the tight pairs.

Theorem 7.1. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote nontrivial pseudo cosine sequences of Γ. Then the following are equivalent.*

(i) $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair.

(ii) There exists a real number ε such that

$$\sigma_i\rho_i - \sigma_{i-1}\rho_{i-1} = \varepsilon(\sigma_{i-1}\rho_i - \sigma_i\rho_{i-1}) \qquad (1 \leq i \leq D). \tag{20}$$

Proof. Set

$$\gamma_i = \sigma_i\rho_i \ (0 \leq i \leq D). \tag{21}$$

(i)$\Longrightarrow$(ii) By Definition 4.1 the sequence $\gamma_0, \gamma_1, \ldots, \gamma_D$ is a pseudo cosine sequence. First assume $\sigma \neq \rho$, and let

$$\varepsilon = \frac{\sigma\rho - 1}{\rho - \sigma}. \tag{22}$$

Pick an integer i $(1 \leq i \leq D)$. By Lemma 3.2, Lemma 3.3 and (22) we have

$$\begin{aligned}
\sigma_i\rho_i - \sigma_{i-1}\rho_{i-1} &= \gamma_i - \gamma_{i-1} \\
&= (\gamma - 1)\,\frac{c_1c_2\cdots c_{i-1}}{b_1b_2\cdots b_{i-1}}\,\sum_{h=0}^{i-1} k_h\gamma_h \\
&= (\sigma\rho - 1)\,\frac{c_1c_2\cdots c_{i-1}}{b_1b_2\cdots b_{i-1}}\,\sum_{h=0}^{i-1} k_h\sigma_h\rho_h \\
&= (\sigma\rho - 1)\,\frac{\sigma_i\rho_{i-1} - \sigma_{i-1}\rho_i}{\sigma - \rho} \\
&= \varepsilon(\sigma_{i-1}\rho_i - \sigma_i\rho_{i-1}),
\end{aligned}$$

as desired. Next assume $\sigma = \rho$. Observe Γ is bipartite or almost bipartite in view of Lemma 6.1. By Theorem 5.3 and since $k\sigma, k\rho$ form a tight pair we find each of $k\sigma, k\rho$ is equal to $-k$. Now by Corollary 5.2 each of σ_i, ρ_i is equal to $(-1)^i$ for $0 \leq i \leq D$. It follows (20) holds for any real number ε.

(ii)$\Longrightarrow$(i) Assume (20) holds for some real number ε. We show $\gamma_0, \gamma_1, \ldots, \gamma_D$ is a pseudo cosine sequence. By Lemma 3.3(i),(iii) it suffices to show

$$(\gamma - 1)\sum_{h=0}^{i} k_h\gamma_h = \frac{b_1b_2\cdots b_i}{c_1c_2\cdots c_i}(\gamma_{i+1} - \gamma_i) \tag{23}$$

for $0 \leq i \leq D - 1$. Pick an integer i $(0 \leq i \leq D - 1)$. By Lemma 3.2 we have

$$(\sigma - \rho)\sum_{h=0}^{i} k_h\sigma_h\rho_h = \frac{b_1b_2\cdots b_i}{c_1c_2\cdots c_i}(\sigma_{i+1}\rho_i - \sigma_i\rho_{i+1}). \tag{24}$$

Setting $i = 1$ in (20) we get

$$\sigma\rho - 1 = \varepsilon(\rho - \sigma). \tag{25}$$

Multiplying equation (24) by ε and simplifying the result using (25), (20) and (21) gives (23) as desired.

□

Definition 7.2. Let Γ denote a distance-regular graph with diameter $D \geq 3$. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. By an *auxiliary parameter* for this pair we mean a real number ε which satisfies Theorem 7.1(ii). We comment on the uniqueness of the auxiliary parameter. Suppose $\sigma = \rho$. Then ε is an auxiliary parameter for all $\varepsilon \in \mathbb{R}$. Suppose $\sigma \neq \rho$. Setting $i = 1$ in (20) we find the auxiliary parameter satisfies

$$\varepsilon = \frac{\sigma\rho - 1}{\rho - \sigma}. \qquad (26)$$

In this case the auxiliary parameter is unique.

8 When can $\sigma_{i-1} = \sigma_i$?

Let Γ denote a distance-regular graph with diameter $D \geq 3$. Assume $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ denote nontrivial pseudo cosine sequences which form a tight pair. Let ε denote a corresponding auxiliary parameter. A bit later in this paper, we discuss certain rational expressions involving $\sigma_0, \sigma_1, \ldots, \sigma_D$ and ε which contain in the denominator the factors $\sigma_{i-1} - \sigma_i$ for $1 \leq i \leq D$. To prepare for this discussion, we investigate how these factors can be zero.

Lemma 8.1. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let us assume this pair has auxiliary parameter $\varepsilon = 1$. Then*

$$(\sigma_{i-1} - \sigma_i)(\rho_{i-1} + \rho_i) = 0 \qquad (1 \leq i \leq D). \qquad (27)$$

Proof. To get (27) set $\varepsilon = 1$ in (20). □

Lemma 8.2. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let us assume Γ is neither bipartite nor almost bipartite. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let us assume this pair has auxiliary parameter $\varepsilon = 1$. Then the following (i)–(iii) hold.*

(i) $\rho_i = (-1)^i$ for $0 \leq i \leq D-1$ and $\rho_D \neq (-1)^D$.

(ii) $\sigma_{D-1} = \sigma_D$.

(iii) $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$.

Proof. (i) Recall $\rho_0 = 1$. Setting $i = 1$ in (27) and recalling $\sigma_0 = 1$, $\sigma \neq 1$ we find $\rho = -1$. Let r denote the maximal integer $(0 \leq r \leq D)$ such that $\rho_i = (-1)^i$ for $0 \leq i \leq r$. We show $r = D-1$. From our preliminary comments $r \geq 1$. Observe $r \neq D$; otherwise Γ is bipartite or almost bipartite in view of Lemma 3.7. Suppose $r \leq D-2$. By the construction $\rho_{r-1} + \rho_r = 0$

and $\rho_r + \rho_{r+1} \neq 0$. Applying Lemma 8.1 we find $\sigma_r = \sigma_{r+1}$. Observe $\sigma_{r+1} \neq \sigma_{r+2}$ by Lemma 3.6 so $\rho_{r+1} + \rho_{r+2} = 0$ by Lemma 8.1. Now $\rho_r + \rho_{r+1} = 0$ by Lemma 3.7, a contradiction. Hence $r = D - 1$.
(ii) Apply (27) with $i = D$ and observe $\rho_{D-1} + \rho_D \neq 0$ by (i) above.
(iii) Recall $a_0 = 0$. We find $a_i = 0$ for $1 \leq i \leq D-2$ by Lemma 3.7 and since $\rho_i = (-1)^i$ for $0 \leq i \leq D-1$. Observe $a_{D-1} \neq 0$; otherwise Γ is bipartite or almost bipartite, contradicting our assumption. □

Lemma 8.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let us assume this pair has auxiliary parameter $\varepsilon = -1$. Then*

$$(\sigma_{i-1} + \sigma_i)(\rho_{i-1} - \rho_i) = 0 \qquad (1 \leq i \leq D). \tag{28}$$

Proof. To get (28) set $\varepsilon = -1$ in (20). □

Lemma 8.4. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let us assume Γ is neither bipartite nor almost bipartite. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let us assume this pair has auxiliary parameter $\varepsilon = -1$. Then the following (i)–(iii) hold.*

(i) $\sigma_i = (-1)^i$ for $0 \leq i \leq D-1$ and $\sigma_D \neq (-1)^D$.

(ii) $\rho_{D-1} = \rho_D$.

(iii) $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$.

Proof. In the proof of Lemma 8.2 interchange the roles of $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$. □

Lemma 8.5. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let ε denote a corresponding auxiliary parameter and assume $\varepsilon \notin \{1, -1\}$. Then $\sigma_{i-1} \neq \sigma_i$ and $\rho_{i-1} \neq \rho_i$ for $1 \leq i \leq D$.*

Proof. First assume $\sigma = \rho$. Observe Γ is bipartite or almost bipartite in view of Lemma 6.1. Applying Theorem 5.3 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ are pseudo cosine sequences for $-k$. By Corollary 5.2 we have $\sigma_i = \rho_i = (-1)^i$ for $0 \leq i \leq D$; so $\sigma_{i-1} \neq \sigma_i$ and $\rho_{i-1} \neq \rho_i$ for $1 \leq i \leq D$ as desired.
Next assume $\sigma \neq \rho$. Then ε is as in (26). Suppose there exists an integer i $(1 \leq i \leq D)$ such that $\sigma_{i-1} = \sigma_i$. Since $\sigma_0 = 1$ and $\sigma \neq 1$ we find $i \geq 2$. Without loss of generality we may assume $\sigma_{i-2} \neq \sigma_{i-1}$. By Lemma 3.5 the common value of σ_{i-1}, σ_i is nonzero. Setting $\sigma_{i-1} = \sigma_i$ in (20) and simplifying we find $(\varepsilon - 1)(\rho_{i-1} - \rho_i) = 0$. We assume $\varepsilon \neq 1$ so $\rho_{i-1} = \rho_i$. From Lemma 3.3(ii) (with i replaced by $i-1$) we find

$$c_{i-1}(\sigma_{i-2} - \sigma_{i-1}) = k(\sigma - 1)\sigma_{i-1}, \tag{29}$$

$$c_{i-1}(\rho_{i-2} - \rho_{i-1}) = k(\rho - 1)\rho_{i-1}. \tag{30}$$

Combining (29), (30) we obtain

$$(\sigma_{i-2}-\sigma_{i-1})(\rho-1)\rho_{i-1}=(\rho_{i-2}-\rho_{i-1})(\sigma-1)\sigma_{i-1}. \tag{31}$$

By (20) (with i replaced by $i-1$) we find

$$\sigma_{i-1}\rho_{i-1}-\sigma_{i-2}\rho_{i-2}=\varepsilon(\sigma_{i-2}\rho_{i-1}-\sigma_{i-1}\rho_{i-2}). \tag{32}$$

Adding $(\sigma-1)\sigma_{i-1}$ times (32) and $\varepsilon\sigma_{i-1}-\sigma_{i-2}$ times (31), and simplifying the result using (26), we routinely find $\sigma_{i-2}=\sigma\sigma_{i-1}$. Evaluating (29) using this we find $c_{i-1}=k$, a contradiction. □

Corollary 8.6. *Let* Γ *denote a distance-regular graph with diameter* $D\geq 3$ *and intersection number* $a_1\neq 0$. *Suppose we are given nontrivial pseudo cosine sequences* $\sigma_0,\sigma_1,\ldots,\sigma_D$ *and* $\rho_0,\rho_1,\ldots,\rho_D$ *which form a tight pair. Let* ε *denote the corresponding auxiliary parameter. Then* $\varepsilon\notin\{1,-1\}$. *Moreover* $\sigma_{i-1}\neq\sigma_i$ *and* $\rho_{i-1}\neq\rho_i$ *for* $1\leq i\leq D$.

Proof. We assume $a_1\neq 0$; therefore Γ is neither bipartite nor almost bipartite. Suppose $\varepsilon=1$. By Lemma 8.2(iii) and since $D\geq 3$ we find $a_1=0$ for a contradiction. Suppose $\varepsilon=-1$. By Lemma 8.4(iii) and since $D\geq 3$ we find $a_1=0$ for a contradiction. Apparently $\varepsilon\notin\{1,-1\}$. Applying Lemma 8.5 we find $\sigma_{i-1}\neq\sigma_i$ and $\rho_{i-1}\neq\rho_i$ for $1\leq i\leq D$. □

9 The case $a_i=0$ $(0\leq i\leq D-2)$, $a_{D-1}\neq 0$.

Lemma 9.1. *Let* Γ *denote a distance-regular graph with diameter* $D\geq 3$ *and valency* k. *Assume* $a_i=0$ *for* $0\leq i\leq D-2$ *and* $a_{D-1}\neq 0$. *Let* $\sigma_0,\sigma_1,\ldots,\sigma_D$ *denote the pseudo cosine sequence for* $-k$. *Then* $\sigma_i=(-1)^i$ *for* $0\leq i\leq D-1$ *and*

$$(-1)^D\sigma_D=1+2a_{D-1}/b_{D-1}. \tag{33}$$

Proof. The sequence $\sigma_0,\sigma_1,\ldots,\sigma_D$ satisfies (4) with $\theta=-k$. It follows $c_i\sigma_{i-1}+b_i\sigma_{i+1}=-k\sigma_i$ for $0\leq i\leq D-2$. Using this and (2), we routinely verify by induction that $\sigma_i=(-1)^i$ for $0\leq i\leq D-1$. To get (33) evaluate (4) at $i=D-1$ and simplify using (2). □

Let Γ denote a distance-regular graph with diameter $D\geq 3$. Assume $a_i=0$ for $0\leq i\leq D-2$ and $a_{D-1}\neq 0$. In what follows we will be concerned with the matrix

$$\begin{pmatrix} 0 & b_0 & & & & & \mathbf{0} \\ c_1 & 0 & b_1 & & & & \\ & c_2 & 0 & \cdot & & & \\ & & \cdot & \cdot & b_{D-3} & & \\ & & & c_{D-2} & 0 & b_{D-2} & \\ \mathbf{0} & & & & c_{D-1} & k-c_{D-1} & \end{pmatrix}. \tag{34}$$

We make a routine observation.

Lemma 9.2. *Let* Γ *denote a distance-regular graph with diameter* $D\geq 3$. *Assume* $a_i=0$ *for* $0\leq i\leq D-2$ *and* $a_{D-1}\neq 0$. *Given scalars* $\sigma_0,\sigma_1,\ldots,\sigma_{D-1},\theta\in\mathbb{R}$, *the following are equivalent.*

- *(i)* *θ is an eigenvalue of the matrix (34), and $(\sigma_0, \sigma_1, \ldots, \sigma_{D-1})^t$ is a corresponding eigenvector, scaled so that $\sigma_0 = 1$.*
- *(ii)* *$\sigma_0, \sigma_1, \ldots, \sigma_{D-1}, \sigma_{D-1}$ is the pseudo cosine sequence for θ.*

Lemma 9.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Assume $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$. Let θ denote any eigenvalue of the matrix (34). Then the following (i)–(iv) hold.*

- *(i)* *$\theta \in \mathbb{R}$.*
- *(ii)* *Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence for θ. Then $\sigma_{D-1} = \sigma_D$.*
- *(iii)* *The pair $\theta, -k$ is tight.*
- *(iv)* *The auxiliary parameter for $\theta, -k$ is 1.*

Proof. (i) Denote the matrix (34) by G. Observe G is tridiagonal. The entries on the superdiagonal and subdiagonal of G are positive. Therefore there exists a diagonal matrix N such that NGN^{-1} is symmetric and has all entries real. We now see the eigenvalues of NGN^{-1} are real. The matrices G and NGN^{-1} have the same eigenvalues, so the eigenvalues of G are real.
(ii) The requirement that $(\sigma_0, \sigma_1, \ldots, \sigma_{D-1})^t$ be an eigenvector of the matrix (34) with $\sigma_0 = 1$ is the same as the requirement that $\sigma_0, \sigma_1, \ldots, \sigma_{D-1}, \sigma_{D-1}$ be the pseudo cosine sequence for θ.
(iii),(iv) Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote the pseudo cosine sequence for θ and let $\rho_0, \rho_1, \ldots, \rho_D$ denote the pseudo cosine sequence for $-k$. Define $\varepsilon = 1$. By Lemma 9.1, we have $\rho_i = (-1)^i$ for $0 \leq i \leq D-1$. By this and since $\sigma_{D-1} = \sigma_D$ we find (20) holds. Applying Theorem 7.1 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Now the pair $\theta, -k$ is tight. By construction the auxiliary parameter for this pair is 1. ☐

Theorem 9.4. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Assume $a_i = 0$ for $0 \leq i \leq D-2$ and $a_{D-1} \neq 0$. Then the following (i)–(iii) hold.*

- *(i)* *The pair θ, k is tight for all $\theta \in \mathbb{R}$.*
- *(ii)* *The pair $\theta, -k$ is tight for any eigenvalue θ of the matrix (34).*
- *(iii)* *Γ has no further tight pairs.*

Proof. (i) This is from Example 4.3.
(ii) Follows from Lemma 9.3(ii).
(iii) Let θ, θ' denote a tight pair, such that $\theta \neq k$, $\theta' \neq k$. Applying Lemma 4.4 we find θ, θ' satisfy (15). By this and since $a_1 = 0$ we find $(\theta + k)(\theta' + k) = 0$. Now $\theta = -k$ or $\theta' = -k$. Without loss of generality we assume $\theta' = -k$. Let ε denote the auxiliary parameter for $\theta, -k$. By (26) (with $\rho = -1$) we find $\varepsilon = 1$. By Lemma 9.2 and Lemma 8.2(ii) we find θ is an eigenvalue of the matrix (34). The result follows. ☐

10 The case $a_1 = 0$ and there exists i $(2 \leq i \leq D-2)$ such that $a_i \neq 0$

Theorem 10.1. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 4$ *and valency* k*. Assume* $a_1 = 0$ *and there exists* i $(2 \leq i \leq D-2)$ *such that* $a_i \neq 0$*.*

(i) The pair θ, k *is tight for all* $\theta \in \mathbb{R}$*.*

(ii) Γ *has no further tight pairs.*

Proof. Let θ, θ' denote a tight pair, such that $\theta \neq k$, $\theta' \neq k$. Applying Lemma 4.4 we find θ, θ' satisfy (15). By this and since $a_1 = 0$ we find $(\theta + k)(\theta' + k) = 0$. Now $\theta = -k$ or $\theta' = -k$. Without loss of generality we assume $\theta' = -k$. Let ε denote the auxiliary parameter for $\theta, -k$. By (26) (with $\rho = -1$) we find $\varepsilon = 1$. By Lemma 8.2(iii) we find $a_i = 0$ for $0 \leq i \leq D-2$, a contradiction. $\square$

11 The case $a_1 \neq 0$

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given nontrivial pseudo cosine sequences $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ which form a tight pair. Let ε denote the corresponding auxiliary parameter. Pick an integer i $(1 \leq i \leq D)$. Note (20) holds; rearranging the terms in (20) we obtain

$$\rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i) = \rho_i(\sigma_i - \varepsilon\sigma_{i-1}). \tag{35}$$

We would like to solve for ρ_{i-1} in (35). However the coefficient $\sigma_{i-1} - \varepsilon\sigma_i$ might be zero. In the following lemma we investigate this possibility.

Lemma 11.1. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$ *and* $a_1 \neq 0$*. Suppose we are given nontrivial pseudo cosine sequences* $\sigma_0, \sigma_1, \ldots, \sigma_D$ *and* $\rho_0, \rho_1, \ldots, \rho_D$ *which form a tight pair. Let* ε *denote the corresponding auxiliary parameter. For* $1 \leq i \leq D-1$ *the following (i)–(iv) are equivalent:*

(i) $\sigma_{i-1} = \varepsilon\sigma_i$*.*

(ii) $\sigma_{i+1} = \varepsilon\sigma_i$*.*

(iii) $\sigma_{i-1} = \sigma_{i+1}$*.*

(iv) $\rho_i = 0$*.*

Proof. Recall line (20) holds.
(i)$\Longrightarrow$ (iv) We replace σ_{i-1} by $\varepsilon\sigma_i$ in (20) to obtain $\sigma_i\rho_i(1-\varepsilon^2) = 0$. Observe $\varepsilon \notin \{1, -1\}$ by Corollary 8.6. Assume for the moment that $\sigma_i = 0$. It follows from the assumption that $\sigma_{i-1} = 0$. So $\sigma_{i-1} = \sigma_i$ contradicting Corollary 8.6. Thus, $\sigma_i \neq 0$ so $\rho_i = 0$.
(iv)$\Longrightarrow$ (i) Setting $\rho_i = 0$ in (20) we obtain $\rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i) = 0$. Observe $\rho_{i-1} \neq 0$ by Lemma 3.5 so $\sigma_{i-1} = \varepsilon\sigma_i$.

(ii)$\Longleftrightarrow$(iv) Similar to the proof of (i)$\Longleftrightarrow$(iv).
(i),(ii)$\Longrightarrow$(iii) Clear.
(iii)$\Longrightarrow$(i) Adding (20) at i and $i+1$ we obtain

$$\sigma_{i+1}\rho_{i+1} - \sigma_{i-1}\rho_{i-1} = \varepsilon(\sigma_i\rho_{i+1} - \sigma_{i+1}\rho_i + \sigma_{i-1}\rho_i - \sigma_i\rho_{i-1}).$$

Replacing σ_{i+1} by σ_{i-1} in the above line and simplifying, we obtain

$$(\sigma_{i-1} - \varepsilon\sigma_i)(\rho_{i+1} - \rho_{i-1}) = 0. \tag{36}$$

We show $\rho_{i+1} - \rho_{i-1} \neq 0$. Suppose $\rho_{i+1} = \rho_{i-1}$. Using this and Lemma 3.6 we have $\rho_{i-1} - \rho_i \neq 0$. Applying Lemma 3.4 to $\rho_0, \rho_1, \dots, \rho_D$ we obtain

$$\frac{a_i}{k} = \frac{\rho_{i-1} - \rho\rho_i}{\rho_{i-1} - \rho_i}. \tag{37}$$

Similarly since $\sigma_{i+1} = \sigma_{i-1}$ we have $\sigma_{i-1} - \sigma_i \neq 0$ and

$$\frac{a_i}{k} = \frac{\sigma_{i-1} - \sigma\sigma_i}{\sigma_{i-1} - \sigma_i}. \tag{38}$$

Combining (37) and (38) we obtain

$$(\sigma_{i-1} - \sigma\sigma_i)(\rho_{i-1} - \rho_i) = (\sigma_{i-1} - \sigma_i)(\rho_{i-1} - \rho\rho_i). \tag{39}$$

We view (20) and (39) as a homogeneous system of linear equations in σ_{i-1} and σ_i. We find that the coefficient matrix of this linear system is given by

$$C = \begin{pmatrix} \varepsilon\rho_i + \rho_{i-1} & -\rho_i - \varepsilon\rho_{i-1} \\ (\rho - 1)\rho_i & \rho_{i-1}(1-\sigma) + \rho_i(\sigma - \rho) \end{pmatrix}.$$

Observe $\sigma \neq \rho$ by Lemma 6.1 so (26) holds. Computing the determinant of C and evaluating the result using (26) we find $\det C = (\rho_{i-1} - \rho_i)(1-\sigma)(\rho_{i-1} - \rho\rho_i)$. We mentioned earlier that $\rho_{i-1} - \rho_i \neq 0$. Observe $1 - \sigma \neq 0$ since the sequence $\sigma_0, \sigma_1, \dots, \sigma_D$ is nontrivial. Observe $\rho_{i-1} - \rho\rho_i \neq 0$; otherwise $a_i = 0$ in view of (37), contradicting Lemma 2.1. We now see that $\det C \neq 0$ so C is nonsingular. Therefore $\sigma_{i-1} = 0$ and $\sigma_i = 0$. This contradicts Lemma 3.5. Thus we have shown $\rho_{i+1} - \rho_{i-1} \neq 0$. By this and (36) we find $\sigma_{i-1} = \varepsilon\sigma_i$. $\square$

Corollary 11.2. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$ *and* $a_1 \neq 0$. *Suppose we are given nontrivial pseudo cosine sequences* $\sigma_0, \sigma_1, \dots, \sigma_D$ *and* $\rho_0, \rho_1, \dots, \rho_D$ *which form a tight pair. Let* ε *denote the corresponding auxiliary parameter. Pick an integer* i $(1 \leq i \leq D-1)$ *and assume the equivalent conditions (i)–(iv) of Lemma 11.1 do not hold. Then both*

$$\rho_{i-1} = \rho_i \frac{\sigma_i - \varepsilon\sigma_{i-1}}{\sigma_{i-1} - \varepsilon\sigma_i}, \tag{40}$$

$$\rho_{i+1} = \rho_i \frac{\sigma_i - \varepsilon\sigma_{i+1}}{\sigma_{i+1} - \varepsilon\sigma_i}. \tag{41}$$

Proof. Observe (20) holds. Rearranging terms we find

$$\rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i) = \rho_i(\sigma_i - \varepsilon\sigma_{i-1}). \tag{42}$$

Observe $\sigma_{i-1} - \varepsilon\sigma_i \neq 0$ by Lemma 11.1(i); solving (42) for ρ_{i-1} we obtain (40). Replacing i by $i+1$ in (42) and rearranging terms we obtain

$$\rho_{i+1}(\sigma_{i+1} - \varepsilon\sigma_i) = \rho_i(\sigma_i - \varepsilon\sigma_{i+1}). \tag{43}$$

Observe $\sigma_{i+1} - \varepsilon\sigma_i \neq 0$ by Lemma 11.1(ii). Solving (43) for ρ_{i+1} we get (41). ☐

12 Tight pseudo cosine sequences

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a nontrivial pseudo cosine sequence. We want to prove there exists at most one nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. To do this we need a lemma.

Lemma 12.1. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and valency k. Suppose there exist real numbers $\theta, \theta', \theta''$ such that (i) none of $\theta, \theta', \theta''$ is equal to k; (ii) $\theta' \neq \theta''$; (iii) the pair θ, θ' is tight; and (iv) the pair θ, θ'' is tight. Then $a_i = 0$ for $0 \leq i \leq D-2$. Moreover $\theta = -k$.*

Proof. By Lemma 4.4 the pair θ, θ' satisfies (15). Applying the same lemma to the pair θ, θ'' and using the fact that $\theta' \neq \theta''$ we obtain $\theta + k/(a_1+1) = 0$. Now in (15) the left-hand side is 0 so the right-hand side is 0. Therefore $a_1 = 0$ and $\theta = -k$. If $D = 3$, we are done. Assume $D \geq 4$. By Theorem 10.1 we have $a_2 = \cdots = a_{D-2} = 0$. The result follows. ☐

Corollary 12.2. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a nontrivial pseudo cosine sequence. Then there exists at most one nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Suppose $\rho_0, \rho_1, \ldots, \rho_D$ exists. Then the corresponding auxiliary parameter is unique.*

Proof. The first assertion follows from Lemma 12.1. The auxiliary parameter is unique by Lemma 6.1 and the comment at the end of Definition 7.2. ☐

Definition 12.3. Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a nontrivial pseudo cosine sequence. We say this sequence is *tight* whenever there exists a nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. By the *auxiliary parameter* for $\sigma_0, \sigma_1, \ldots, \sigma_D$ we mean the auxiliary parameter for the tight pair $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$.

Lemma 12.4. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given a nontrivial pseudo cosine sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ which is tight in the sense of Definition 12.3. Let ε denote the corresponding auxiliary parameter. Then the following (i)–(iv) hold.*

(i) $\sigma_2 \neq 1$.

(ii) $\varepsilon\sigma \neq 1$.

(iii) $\sigma_2 \neq \varepsilon\sigma$.

(iv) $\sigma_2 \neq \sigma^2$.

Proof. By Definition 12.3 there exists a nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Thus (20) holds.
(i) Setting $i = 3$ in (20) we find

$$\sigma_3\rho_3 - \sigma_2\rho_2 - \varepsilon(\sigma_2\rho_3 - \sigma_3\rho_2) \tag{44}$$

is zero. We assume $\sigma_2 = 1$ and show (44) is not zero. In order to do this we evaluate the terms in (44). We assume $\sigma_2 = 1$; therefore the equivalent conditions (i)–(iv) hold in Lemma 11.1 for $i = 1$. Setting $i = 1$ in Lemma 11.1(iv) we find $\rho = 0$. Similarly using Lemma 11.1(i) we find $\varepsilon\sigma = 1$. It follows that $\sigma \neq 0$ and $\varepsilon = 1/\sigma$. Setting $i = 1$ in (13), solving for σ and eliminating a_1 in the result using (2) we get

$$\sigma = -\frac{1 + b_1}{k}. \tag{45}$$

By this and since $\varepsilon = 1/\sigma$ we find

$$\varepsilon = -\frac{k}{1 + b_1}. \tag{46}$$

Setting $i = 2$ in (4), solving for σ_3 and eliminating c_2, σ in the result using (2), (45) we get

$$\sigma_3 = -\frac{(a_2 + b_2)(1 + b_1) + ka_2}{kb_2}. \tag{47}$$

Applying (4) to the sequence $\rho_0, \rho_1, \ldots, \rho_D$ we find

$$c_i\rho_{i-1} + a_i\rho_i + b_i\rho_{i+1} = k\rho\rho_i \quad (0 \leq i \leq D - 1). \tag{48}$$

Setting $i = 1, 2$ in (48) and solving for ρ_2, ρ_3 respectively using $\rho = 0$ we obtain

$$\rho_2 = -{b_1}^{-1}, \tag{49}$$

$$\rho_3 = a_2{b_1}^{-1}{b_2}^{-1}. \tag{50}$$

Evaluating (44) using (45), (46), (47) and (49), (50) we routinely verify that

$$\sigma_3\rho_3 - \sigma_2\rho_2 - \varepsilon(\sigma_2\rho_3 - \sigma_3\rho_2) = -a_2\frac{(a_2 + b_2)(1 + b_1 + k)}{b_1 b_2^2 k}. \tag{51}$$

Recall the left-hand side of (51) is 0 so the right-hand side is 0. It follows $a_2 = 0$. However $a_2 \neq 0$ by Lemma 2.1 and since $a_1 \neq 0$. We have now shown $\sigma_2 \neq 1$.
(ii),(iii) Immediate from (i) above and Lemma 11.1.
(iv) Suppose $\sigma_2 = \sigma^2$. Setting $i = 1$ in (13) and since $\sigma \neq 1$ we find $\sigma = -1/(a_1 + 1)$. Setting $\theta = k\sigma, \theta' = k\rho$ in equation (15) we find that the left-hand side is 0 while the right-hand side is nonzero. This violates Lemma 4.4. Thus $\sigma_2 \neq \sigma^2$. □

Lemma 12.5. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$ *and* $a_1 \neq 0$*. Suppose we are given nontrivial pseudo cosine sequences* $\sigma_0, \sigma_1, \ldots, \sigma_D$ *and* $\rho_0, \rho_1, \ldots, \rho_D$ *which form a tight pair. Let* ε *denote the corresponding auxiliary parameter. Then the following (i)–(iii) hold.*

(i) $\sigma \neq \varepsilon$.

(ii) $\rho = \dfrac{1-\varepsilon\sigma}{\sigma-\varepsilon}$.

(iii) $\rho_2 = \dfrac{\rho(\sigma-\varepsilon\sigma_2)}{\sigma_2-\varepsilon\sigma}$.

Proof. (i) Suppose $\sigma = \varepsilon$. Then setting $i = 1$ in (20) we find $\varepsilon^2 = 1$; contradicting Corollary 8.6.
(ii) Solve (26) for ρ.
(iii) Set $i = 1$ in (41). ☐

13 The intersection numbers

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given a nontrivial pseudo cosine sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ which is tight. Let ε denote the corresponding auxiliary parameter. In this section we compute the intersection numbers in terms of $\sigma_0, \sigma_1, \ldots, \sigma_D$ and ε.

We begin with the valency k.

Lemma 13.1. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$ *and* $a_1 \neq 0$*. Suppose we are given a nontrivial pseudo cosine sequence* $\sigma_0, \sigma_1, \ldots, \sigma_D$ *which is tight. Let* ε *denote the corresponding auxiliary parameter. Then the valency* k *satisfies*

$$k = h\frac{\sigma-\varepsilon}{\sigma-1} \tag{52}$$

where

$$h = \frac{(1-\sigma)(1-\sigma_2)}{(\sigma^2-\sigma_2)(1-\varepsilon\sigma)}. \tag{53}$$

We remark the denominator in (52) is nonzero since $\sigma_0, \sigma_1, \ldots, \sigma_D$ *is nontrivial. Moreover the denominator in (53) is nonzero by Lemma 12.4.*

Proof. By Definition 12.3 there exists a nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Since $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a pseudo cosine sequence we find it satisfies (8). Setting $i = 1$ in (8) we get

$$k(\sigma^2-\sigma) + b_1(\sigma-\sigma_2) = 1-\sigma. \tag{54}$$

Similarly

$$k(\rho^2-\rho) + b_1(\rho-\rho_2) = 1-\rho. \tag{55}$$

We view (54) and (55) as a system of linear equations in k and b_1. The coefficient matrix is

$$E = \begin{pmatrix} \sigma^2 - \sigma & \sigma - \sigma_2 \\ \rho^2 - \rho & \rho - \rho_2 \end{pmatrix}.$$

Evaluating the determinant of E using Lemma 12.5(ii),(iii) we find that

$$\det E = \frac{(1+\varepsilon)(1-\sigma)(1-\varepsilon\sigma)(\sigma-\sigma_2)(\sigma^2-\sigma_2)}{(\sigma-\varepsilon)^2(\sigma_2-\varepsilon\sigma)}.$$

We show $\det E \neq 0$. Observe $1+\varepsilon \neq 0$ by Corollary 8.6, $1-\sigma \neq 0$ since the sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ is nontrivial, $1-\varepsilon\sigma \neq 0$ by Lemma 12.4(ii), $\sigma - \sigma_2 \neq 0$ by Corollary 8.6 and $\sigma^2 - \sigma_2 \neq 0$ by Lemma 12.4(iv). We have now shown $\det E \neq 0$. Solving the system (54), (55) we find

$$k = -\frac{(\sigma-\varepsilon)(1-\sigma_2)}{(\sigma^2-\sigma_2)(1-\varepsilon\sigma)}.$$

The desired result follows. ☐

Lemma 13.2. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given a nontrivial pseudo cosine sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ which is tight. Let ε denote the corresponding auxiliary parameter. Then*

$$a_i = g\frac{(\sigma_{i+1}-\sigma\sigma_i)(\sigma_{i-1}-\sigma\sigma_i)}{(\sigma_{i+1}-\sigma_i)(\sigma_{i-1}-\sigma_i)} \qquad (1 \leq i \leq D-1) \tag{56}$$

where

$$g = \frac{(\varepsilon-1)(1-\sigma_2)}{(\sigma^2-\sigma_2)(1-\varepsilon\sigma)}. \tag{57}$$

We remark the denominator in (56) is nonzero by Corollary 8.6. Moreover the denominator in (57) is nonzero by Lemma 12.4.

Proof. By Definition 12.3 there exists a nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Pick an integer i $(1 \leq i \leq D-1)$. By Lemma 3.4 and since $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a pseudo cosine sequence we find

$$c_i(\sigma_{i+1}-\sigma_{i-1}) + a_i(\sigma_{i+1}-\sigma_i) = k(\sigma_{i+1}-\sigma\sigma_i). \tag{58}$$

Similarly

$$c_i(\rho_{i+1}-\rho_{i-1}) + a_i(\rho_{i+1}-\rho_i) = k(\rho_{i+1}-\rho\rho_i). \tag{59}$$

We solve (58), (59) for a_i. We consider two cases. First assume $\sigma_{i+1} - \sigma_{i-1} \neq 0$. We view (58) and (59) as a system of linear equations in c_i and a_i. The coefficient matrix of the above system is given by

$$F = \begin{pmatrix} \sigma_{i+1}-\sigma_{i-1} & \sigma_{i+1}-\sigma_i \\ \rho_{i+1}-\rho_{i-1} & \rho_{i+1}-\rho_i \end{pmatrix}.$$

We evaluate the determinant of F. We assume $\sigma_{i+1} \neq \sigma_{i-1}$ so Corollary 11.2 applies; therefore ρ_{i-1}, ρ_{i+1} in F are given by (40), (41) respectively. Eliminating ρ_{i-1} and ρ_{i+1} in F using (40) and (41) we get

$$\det F = \rho_i \frac{(1+\varepsilon)(\sigma_{i-1}-\sigma_i)(\sigma_{i+1}-\sigma_i)(\sigma_{i-1}-\sigma_{i+1})}{(\sigma_{i-1}-\varepsilon\sigma_i)(\sigma_{i+1}-\varepsilon\sigma_i)}.$$

We show $\det F \neq 0$. Observe $\rho_i \neq 0$ by Lemma 11.1 and since $\sigma_{i+1} \neq \sigma_{i-1}$. Observe $1+\varepsilon \neq 0$, $\sigma_{i-1}-\sigma_i \neq 0$ and $\sigma_{i+1}-\sigma_i \neq 0$ by Corollary 8.6. Now $\det F \neq 0$. Solving (58), (59) for a_i and then evaluating the result using Corollary 11.2, Lemma 12.5(ii) and Lemma 13.1 we routinely obtain (56), (57). We have now proved the result for the case $\sigma_{i+1}-\sigma_{i-1} \neq 0$. Next assume $\sigma_{i+1}-\sigma_{i-1} = 0$. Observe $\sigma_{i+1}-\sigma_i \neq 0$ by Lemma 3.6. Solving for a_i in (58) we get

$$a_i = k\frac{\sigma_{i+1}-\sigma\sigma_i}{\sigma_{i+1}-\sigma_i}. \tag{60}$$

Eliminating k in (60) using (52) and (53) we get

$$a_i = \frac{(1-\sigma_2)(\varepsilon-\sigma)(\sigma_{i+1}-\sigma\sigma_i)}{(\sigma^2-\sigma_2)(1-\varepsilon\sigma)(\sigma_{i+1}-\sigma_i)}. \tag{61}$$

We eliminate $\varepsilon - \sigma$ in (61). By Lemma 11.1 and since $\sigma_{i+1} = \sigma_{i-1}$ we find $\sigma_{i-1} = \varepsilon\sigma_i$. Observe $\sigma_{i-1}-\sigma_i \neq 0$ by Corollary 8.6. Using these facts we find

$$\varepsilon - \sigma = \frac{(\varepsilon-1)(\sigma_{i-1}-\sigma\sigma_i)}{\sigma_{i-1}-\sigma_i}. \tag{62}$$

Eliminating $\varepsilon - \sigma$ in (61) using (62) we get (56), (57). $\square$

Lemma 13.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given a nontrivial pseudo cosine sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ which is tight. Let ε denote the corresponding auxiliary parameter. Then*

$$b_i(\sigma_{i-1}-\sigma_{i+1}) = h\frac{(\sigma_{i-1}-\sigma\sigma_i)(\sigma_{i+1}-\varepsilon\sigma_i)}{\sigma_{i+1}-\sigma_i} \qquad (1 \leq i \leq D-1), \tag{63}$$

$$c_i(\sigma_{i+1}-\sigma_{i-1}) = h\frac{(\sigma_{i+1}-\sigma\sigma_i)(\sigma_{i-1}-\varepsilon\sigma_i)}{\sigma_{i-1}-\sigma_i} \qquad (1 \leq i \leq D-1), \tag{64}$$

where h is as in (53). We remark the denominators in (63), (64) are nonzero by Corollary 8.6.

Proof. To obtain (63), in line (12) evaluate k, a_i using Lemmas 13.1, 13.2 respectively. Line (64) is similarly obtained. $\square$

14 A characterization theorem

Definition 14.1. Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ and ε denote real numbers such that $\sigma_0 = 1$. We consider several conditions on these scalars.

(i) By condition A we mean $\varepsilon \neq -1$, equations (56), (57) hold and the denominators in (56), (57) are nonzero.

(ii) By condition B we mean equations (63), (53) hold and the denominators in (63), (53) are nonzero.

(iii) By condition C we mean equations (52), (64), (53) hold and the denominators in (52), (64),(53) are nonzero.

Theorem 14.2. *Let* Γ *denote a distance-regular graph with diameter* $D \geq 3$ *and* $a_1 \neq 0$. *Let* $\sigma_0, \sigma_1, \ldots, \sigma_D$ *and* ε *denote real numbers such that* $\sigma_0 = 1$. *Then with reference to Definition 14.1 the following (i)–(iv) are equivalent.*

(i) $\sigma_0, \sigma_1, \ldots, \sigma_D$ *is a tight nontrivial pseudo cosine sequence and* ε *is the corresponding auxiliary parameter.*

(ii) $\sigma_0, \sigma_1, \ldots, \sigma_D$ *is a nontrivial pseudo cosine sequence and* $\sigma_0, \sigma_1, \ldots, \sigma_D, \varepsilon$ *satisfy* A.

(iii) $\sigma_0, \sigma_1, \ldots, \sigma_D, \varepsilon$ *satisfy both* A *and* B.

(iv) $\sigma_0, \sigma_1, \ldots, \sigma_D, \varepsilon$ *satisfy both* A *and* C.

Proof. (i)$\Longrightarrow$(ii) Applying Definition 12.3 there exists a nontrivial pseudo cosine sequence $\rho_0, \rho_1, \ldots, \rho_D$ such that $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Applying Corollary 8.6 to this pair we find $\varepsilon \neq -1$. The result follows by Lemma 13.2.
(ii) $\Longrightarrow$ (iii) We verify that $\sigma_0, \sigma_1, \ldots, \sigma_D, \varepsilon$ satisfy condition B. In each of (63), (53) the denominator is nonzero because each factor in this denominator is in the denominator of (56), (57). We now verify (63), (53). Observe Lemma 3.4(i) holds. Applying that lemma we find (12), (13) hold. Setting $i = 1$ in (13) we find $k(\sigma^2 - \sigma_2) = 1 - \sigma_2 + a_1(\sigma - \sigma_2)$. Observe $\sigma^2 - \sigma_2$ is not zero since it is a factor in the denominator of (57). Solving the equation for k and simplifying the result using (56) we obtain (52), (53). Evaluating (12) using this and (56) we obtain (63). The result follows.
(iii) $\Longrightarrow$ (iv) We verify that $\sigma_0, \sigma_1, \ldots, \sigma_D, \varepsilon$ satisfy condition C. The denominator in (53) is nonzero by condition B. In each of (52), (64) the denominator is nonzero because each factor in this denominator is in the denominator of (53), (56), (57), (63). We now verify (52), (64), (53). In order to do this we first obtain (52). Observe that in (63) the coefficient of b_1 is equal to $1 - \sigma_2$. This coefficient is not zero; if it is then $g = 0$ in view of (57) which implies $a_1 = 0$ for a contradiction. Now to obtain (52), simplify the right-hand side of $k = b_1 + a_1 + 1$ using (56), (63),(57), (53). We now have (52). To obtain (64), expand the left-hand side using (2) and simplify the result using (56), (52), (63).
(iv) $\Longrightarrow$ (ii) We show $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a nontrivial pseudo cosine sequence. To do this we apply Lemma 3.4(i),(iii). Using (56), (52), (64) and the fact that $g = h\frac{\varepsilon - 1}{1 - \sigma}$ from (53) and (57) we routinely verify (13). Observe Lemma 3.4(iii) holds. Applying that lemma we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ is a pseudo cosine sequence. We remark $\sigma \neq 1$ since $1 - \sigma$ is in the denominator of (56). Therefore $\sigma_0, \sigma_1, \ldots, \sigma_D$ is nontrivial.
(ii), (iii), (iv) $\Longrightarrow$ (i) Observe $\sigma \neq 1$ since $\sigma_0, \sigma_1, \ldots, \sigma_D$ is nontrivial. Observe $\sigma \neq \varepsilon$ by (52) and since $k \neq 0$. Define

$$\rho = \frac{1 - \varepsilon\sigma}{\sigma - \varepsilon}. \tag{65}$$

Let $\rho_0, \rho_1, \ldots, \rho_D$ denote the pseudo cosine sequence for $k\rho$. Observe $\rho \neq 1$; otherwise $\sigma = 1$ or $\varepsilon = -1$ for a contradiction. Therefore $\rho_0, \rho_1, \ldots, \rho_D$ is nontrivial. We show $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. To do this we apply Theorem 7.1. Specifically we show that for any integer i $(1 \leq i \leq D)$

$$\rho_i(\sigma_i - \varepsilon\sigma_{i-1}) = \rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i). \tag{66}$$

We do this by induction on i. Observe that when $i = 1$ (66) holds in view of (65). Now pick an integer i $(1 \leq i \leq D-1)$ and assume (66) holds. We show that $\rho_{i+1}(\sigma_{i+1} - \varepsilon\sigma_i) = \rho_i(\sigma_i - \varepsilon\sigma_{i+1})$. This will follow provided we can show

$$b_i\rho_{i+1}(\sigma_{i+1} - \varepsilon\sigma_i) - b_i\rho_i(\sigma_i - \varepsilon\sigma_{i+1}) = 0. \tag{67}$$

We evaluate the expression on the left-hand side of (67) in the following manner. In that expression, we eliminate the first occurrence of b_i using (4) and we eliminate the second occurrence of b_i using (2). We conclude the left-hand side of (67) is equal to $x + y + z$ where

$$\begin{aligned} x &= k\rho\rho_i(\sigma_{i+1} - \varepsilon\sigma_i) - k\rho_i(\sigma_i - \varepsilon\sigma_{i+1}) - a_i\rho_i(\sigma_{i+1} - \sigma_i)(1+\varepsilon), && (68)\\ y &= -c_i\rho_{i-1}(\sigma_{i+1} - \sigma_{i-1}), && (69)\\ z &= c_i\rho_i(\sigma_i - \varepsilon\sigma_{i+1}) - c_i\rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i). && (70) \end{aligned}$$

Eliminating $\rho_{i-1}(\sigma_{i-1} - \varepsilon\sigma_i)$ in (70) using (66) we get

$$y + z = -c_i(\sigma_{i+1} - \sigma_{i-1})(\rho_{i-1} + \varepsilon\rho_i). \tag{71}$$

We now simplify $x+y+z$. In (68) we eliminate a_i, k, ρ using (56),(52), (65) and simplify the result using the fact that $g = h\frac{\varepsilon-1}{1-\sigma}$ from (57), (53); and in (71) we eliminate $c_i(\sigma_{i+1} - \sigma_{i-1})$ using (64). We routinely obtain $x+y+z$ is equal to $h\frac{\sigma_{i+1}-\sigma\sigma_i}{\sigma_{i-1}-\sigma_i}$ times Δ where Δ is the left-hand side of (66) minus the right-hand side of (66). Observe $\Delta = 0$ by (66) so $x+y+z = 0$. Thus we have (67). We have now shown by induction that (66) holds for $1 \leq i \leq D$. Applying Theorem 7.1 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ and $\rho_0, \rho_1, \ldots, \rho_D$ form a tight pair. Comparing (26), (65) we find ε is the corresponding auxiliary parameter. Applying Definition 12.3 we find $\sigma_0, \sigma_1, \ldots, \sigma_D$ is tight and ε is the corresponding auxiliary parameter. $\square$

15 Feasible pseudo cosine sequences

Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Suppose we are given a nontrivial pseudo cosine sequence $\sigma_0, \sigma_1, \ldots, \sigma_D$ which is tight. In view of Lemma 13.3 it is natural to consider the case in which $\sigma_{i-1} \neq \sigma_{i+1}$ for $1 \leq i \leq D-1$. We now investigate this case.

Definition 15.1. Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \ldots, \sigma_D$ denote a pseudo cosine sequence. We say this sequence is *feasible* whenever the following (i),(ii) hold.

(i) $\sigma_{i-1} \neq \sigma_{i+1}$ for $1 \leq i \leq D-1$,

(ii) $\sigma_0, \sigma_1, \dots, \sigma_D$ is tight.

Theorem 15.2. *Let Γ denote a distance-regular graph with diameter $D \geq 3$. Let $\sigma_0, \sigma_1, \dots, \sigma_D$ and ε, h denote real numbers. Then the following (i), (ii) are equivalent.*

(i) $a_1 \neq 0$; the sequence $\sigma_0, \sigma_1, \dots, \sigma_D$ is a feasible pseudo cosine sequence of Γ, ε is the corresponding auxiliary parameter, and

$$h = \frac{(1-\sigma)(1-\sigma_2)}{(\sigma^2 - \sigma_2)(1-\varepsilon\sigma)}. \tag{72}$$

(ii) $\sigma_0 = 1$, $\varepsilon \neq 1$, $\varepsilon \neq -1$,

$$\begin{aligned} k &= h\frac{\sigma - \varepsilon}{\sigma - 1}, & & (73)\\ b_i &= h\frac{(\sigma_{i-1} - \sigma\sigma_i)(\sigma_{i+1} - \varepsilon\sigma_i)}{(\sigma_{i-1} - \sigma_{i+1})(\sigma_{i+1} - \sigma_i)} \qquad (1 \leq i \leq D-1), & & (74)\\ c_i &= h\frac{(\sigma_{i+1} - \sigma\sigma_i)(\sigma_{i-1} - \varepsilon\sigma_i)}{(\sigma_{i+1} - \sigma_{i-1})(\sigma_{i-1} - \sigma_i)} \qquad (1 \leq i \leq D-1), & & (75) \end{aligned}$$

and the denominators in (73)–(75) are nonzero.

Proof. (i) $\Longrightarrow$ (ii) Observe $\sigma_0 = 1$ by Lemma 3.4 and since $\sigma_0, \sigma_1, \dots, \sigma_D$ is a pseudo cosine sequence. By Corollary 8.6 we have $\varepsilon \notin \{1, -1\}$. The remaining assertions follow from Theorem 14.2.
(ii) $\Longrightarrow$ (i) First we show (72). Setting $i = 1$ in (75) and solving for h using the fact that $c_1 = 1$ we get (72). Observe $h \neq 0$; otherwise $k = 0$. Next we show that $\sigma_0, \sigma_1, \dots, \sigma_D$ is a pseudo cosine sequence. To do this we apply Lemma 3.3(i),(ii). Using (73), (74), (75) we routinely verify that (8) holds. By Lemma 3.3 $\sigma_0, \sigma_1, \dots, \sigma_D$ is a pseudo cosine sequence. This sequence is nontrivial; otherwise $h = 0$. Solving for a_i using (2) and (74), (75) we routinely verify that (56), (57) hold. We show $a_1 \neq 0$. Setting $i = 1$ in (56) we obtain

$$a_1 = \frac{(1-\varepsilon)(1+\sigma)(1-\sigma_2)}{(1-\varepsilon\sigma)(\sigma_2 - \sigma)}. \tag{76}$$

Observe $1 - \varepsilon \neq 0$ by assumption; $1 + \sigma \neq 0$ by (74) and the fact that $b_1 \neq 0$; and $1 - \sigma_2 \neq 0$ otherwise $h = 0$. It follows $a_1 \neq 0$. Applying Definition 14.1(i) we find $\sigma_0, \sigma_1, \dots, \sigma_D$ and ε together satisfy condition A. Now by Theorem 14.2(i),(ii) we find $\sigma_0, \sigma_1, \dots, \sigma_D$ is tight and that ε is the corresponding auxiliary parameter. We observe $\sigma_{i-1} \neq \sigma_{i+1}$ $(1 \leq i \leq D-1)$ since the denominators in (74) are nonzero. Now $\sigma_0, \sigma_1, \dots, \sigma_D$ is feasible by Definition 15.1. ☐

We finish this section with a comment.

Lemma 15.3. *Let Γ denote a distance-regular graph with diameter $D \geq 3$ and $a_1 \neq 0$. Let $\sigma_0, \sigma_1, \dots, \sigma_D$ denote a feasible pseudo cosine sequence. Let $\rho_0, \rho_1, \dots, \rho_D$ denote a nontrivial* *pseudo cosine sequence such that* $\sigma_0, \sigma_1, \ldots, \sigma_D$ *and* $\rho_0, \rho_1, \ldots, \rho_D$ *form a tight pair. Let* ε *denote the corresponding auxiliary parameter. Then*

$$\rho_i = \prod_{j=1}^{i} \frac{\sigma_{j-1} - \varepsilon\sigma_j}{\sigma_j - \varepsilon\sigma_{j-1}} \qquad (0 \leq i \leq D) \tag{77}$$

and the denominators in (77) are all nonzero.

Proof. We first show the denominators in (77) are all nonzero. By Lemma 11.1 and since $\sigma_{i-1} - \sigma_{i+1} \neq 0$ for $1 \leq i \leq D-1$ we find $\sigma_i - \varepsilon\sigma_{i-1} \neq 0$ for $2 \leq i \leq D$. Observe $\sigma \neq \varepsilon$ in view of Lemma 12.5. Using this and the fact $\sigma_0 = 1$ we get $\sigma_i - \varepsilon\sigma_{i-1} \neq 0$ for $1 \leq i \leq D$. We have now shown the denominators in (77) are all nonzero. We now verify (77). Observe (77) holds for $i = 0$ since $\rho_0 = 1$. Line (77) holds for $i = 1$ by (26). Line (77) holds for $2 \leq i \leq D$ by (41) and a routine induction. $\square$

References

[1] E. Bannai and T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin-Cummings Lecture Note Ser. 58, Benjamin-Cummings, Menlo Park, CA 1984.

[2] A.E. Brouwer, A.M. Cohen and A. Neumaier, *Distance-Regular Graphs*, Springer, New York, 1989.

[3] J. T. Go and P. Terwilliger, Tight distance-regular graphs and the subconstituent algebra, European J. Combin., 23 (2002), 793–816.

[4] A. Jurišić, AT4 family and 2-homogeneous graphs, Preprint.

[5] A. Jurišić and J. Koolen, A local approach to 1-homogeneous graphs, Des. Codes Cryptography, 21 (2000), 127–147.

[6] A. Jurišić and J. Koolen, Nonexistence of some antipodal distance-regular graphs of diameter four, European J. Combin., 21 (2000), 1039–1046.

[7] A. Jurišić and J. Koolen, 1-homogeneous graphs with cocktail party μ-graphs, J. Algebraic Combin., To appear.

[8] A. Jurišić and J. Koolen, Krein parameters and antipodal distance-regular graphs with diameter 3 and 4, Discrete Math., 244 (2002), 181–202.

[9] A. Jurišić, J. Koolen and P. Terwilliger, Tight distance-regular graphs, J. Algebraic Combin., 12 (2000), 163–197.

[10] A. Jurišić and P. Terwilliger, Pseudo tight distance-regular graphs, Preprint.

[11] M. MacLean, An inequality involving two eigenvalues of a bipartite distance-regular graph, Discrete Math., 225 (2000), 193–216.

[12] M. MacLean, Taut distance-regular graphs of odd diameter, J. Algebraic Combin., Submitted.

[13] M. MacLean, Taut distance-regular graphs of even diameter, J. Combin. Theory Ser. B, Submitted.

[14] A. A. Pascasio, Tight graphs and their primitive idempotents, J. Algebraic Combin., 10 (1999), 47-59.

[15] A. A. Pascasio, Tight distance-regular graphs and Q-polynomial property, Graphs and Combin., 17 (2001), 149–169.

[16] A. A. Pascasio, An inequality on the cosines of a tight distance-regular graph, Linear Algebra Appl., 325 (2001), 147–159.

[17] A. A. Pascasio, An inequality in character algebras, Discrete Math., 264 (2003), 201–209.

[18] M. Tomiyama, On the primitive idempotents of distance-regular graphs, Discrete Math., 240 (2001), 281–294.

[19] C.W. Weng and P. Terwilliger, Distance-regular graphs, pseudo primitive idempotents, and the Terwilliger algebra, Seidel 80 Conference, To appear.

Arlene A. Pascasio
Department of Mathematics
De La Salle University - Manila
2401 Taft Avenue
Malate Manila 1004
Philippines
Email: pascasioa@dlsu.edu.ph

Paul Terwilliger
Department of Mathematics
University of Wisconsin - Madison
Van Vleck Hall
480 Lincoln Drive
Madison WI, USA 53706-1388
Email: terwilli@math.wisc.edu